

AD-A205 432

REPORT DOCUMENTATION PAGE				
1a. REPORT SECURITY CLASSIFICATION <i>Unclassified</i>		1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		7a. NAME OF MONITORING ORGANIZATION European Office of Aerospace Research and Development		
5a. NAME OF PERFORMING ORGANIZATION		6b. OFFICE SYMBOL (if applicable)		7b. ADDRESS (City, State, and ZIP Code) Box 14 FPO New York 09510-0200
6a. ADDRESS (City, State, and ZIP Code) The Auroral Observatory P.O. Box 953 N-9001 Tromsø, Norway		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
7a. NAME OF FUNDING/SPONSORING ORGANIZATION EOARD		8b. OFFICE SYMBOL (if applicable)		10. SOURCE OF FUNDING NUMBERS
8c. ADDRESS (City, State, and ZIP Code)		PROGRAM ELEMENT NO 61102F	PROJECT NO 2301	TASK NO D1
11. TITLE (Include Security Classification) Optical Studies of the Magnetospheric Cusp from Svalbard				
12. PERSONAL AUTHOR(S) Kjell Henriksen				
13a. TYPE OF REPORT	13b. TIME COVERED FROM 30.9-87 TO 30.9.88	14. DATE OF REPORT (Year, Month, Day) 89-1-2	15. PAGE COUNT 3	
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)				
<p>The present investigation was made on spectrometric auroral data obtained on the dayside and nightside of the auroral oval in Svalbard. Synthetic spectra of $O_2^+ 1NG$ and $N_2^+ 1NG$ bands were calculated. The transition probabilities were obtained by particular solutions of the Schrödinger equation. The synthetic spectra were used to analyse the auroral spectrum from 5200 to 5800 Å, deriving the intensities of all the auroral emissions of this spectral region.</p> <p>The same spectral technique was used to analyse solar spectra, deriving ozone column densities and spectral UV irradiances. <i>Keywords: Ultraviolet spectrometry; Magnetosphere; Arctic Ocean; Auroras; Europe; Spectrometry; etc.</i></p> <p>(N2+)</p>				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Owen R. Cote		22b. TELEPHONE (Include Area Code)		22c. OFFICE SYMBOL EOARD/LDA

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.
All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

89 3 06 142

This report has been reviewed and is releasable to the National Technical Information Service (NTIS). At NTIS it will be releasable to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

On 12-6-68

OWEN R. COTE'
Chief, Chief, Geophysics and Space

Fred T. Gilliam

FRED T. GILLIAM, Lt Col, USAF
Chief Scientist



Letter to file

A-1

Final Scientific Report

30 Sep 87 - 29 Sep 88

By the AFOSR-86-0327 grant we have been able to analyze our optical data from Svalbard. We have shown that the optical spectra can be precisely reproduced by synthetic spectra, where the transition probabilities have been obtained from particular solutions of the Schrödinger equation.

This method has been used on measurements of the N_2^+ and O_2^+ first negative bands, absolute temperatures and intensity ratios. Neutral temperature variations in the auroral E-region are difficult to obtain, but we have published an able method. Since both the O_2^+ and N_2^+ mainly originate in the E-region, the rotational distribution of these bands will indicate temperature variations. Intensity ratios of these band systems will point out height variations of the centroid of the emission layer since the density of N_2 and O_2 varies through the E-layer. However, a better estimate of the height could be obtained by triangulation measurements using two ground-based instruments situated about 100 km apart along a geomagnetic meridian. Then the whole auroral height profile would be found.

Curious enough, no optical measurements have ever shown a temperature increase of the neutral atmosphere in the E-region associated with an auroral substorm. The same result is obtained in our work, and we argue that it is due to the high heat capacity of the E-region and rapid transport processes.

The findings of this analysis are published in two papers:

K. Henriksen and L. Veseth: Analysis of auroral O_2^+ first negative bands. Can. J. Phys. 65, 1119-1132, 1987.

K. Henriksen, L. Veseth, C.S. Deehr and R.W. Smith: Neutral temperatures and emission height changes in an E-region aurora. Planet. Space Sci. 35, 1317-1321, 1987.

During the auroral season 1986/87 cooperative studies with Dr. W. Pendelton's group, Utah State University, were carried out

in Svalbard. They measured with a Michelson Interferometer the spectral region from 9000 to 15000 Å. In this spectral region the N_2^+ Meinel (0,0) band and the distinguished HeI line 10830 Å appear. Preliminary results indicate that the N_2^+ M (0,0) band is selectively enhanced in F-region aurora by energy transfer from O^+ ions. The Meinel bands which we measure by spectrometer in the near infrared, are not enhanced. This phenomenon is detectable only in cusp and polar cap aurora.

Helium precipitation seems to occur intermittently in bright auroras, and the intensity enhancement of the 10830 Å line can be 2-3 times the geocoronal background.

During the period 1987-88 Dr. C.S. Gardner, University of Illinois at Urbana - Champaign, had a dimer laser in Svalbard to study the structure of the mesospheric sodium layer, gravity waves and how they were affected by geomagnetic and auroral activity. Primary results indicate that the sodium layer almost disappears during summer months, but influence from geomagnetic activity is harder to see. Analysis is under way.

The analysis programs developed under this grant will be used in future auroral and optical cooperative projects with American groups visiting Svalbard. Our station, Nordlysstasjonen, will be extended by funds from National Science Foundation, and it is proposed to be a main station for the CEDAR program.

The spectrometric analysis technique is also usable for studies of solar irradiance and ozone. We have developed spectrometric instruments capable of measuring the solar irradiance from 2800 to 8000 Å, and used it in Tromsø and Svalbard. The media and news are filled with predictions and hints about decreasing stratospheric ozone and increasing UVB irradiance. Here we have developed a method for simultaneously measuring both UV irradiance and the ozone column density. We find that both quantities have the expected values, and we don't find any published measurement showing that the global ozone layer is in danger. However, the frightening theories are supported by Nimbus 7 SBUV measurement, but they are wrong due to the decreasing sensitivity of the SBUV instrument. Our first results are accepted for publication in Atmospheric Environment:

K. Henriksen, K. Stamnes and P. Østensen: Measurements of solar UV, visible and near IR irradiance at 78⁰N, 1988.

Manuscript included. The referees comment that the manuscript gives the first results in a very important measurement program, and we will of course continue.

The Auroral Observatory

2 Jan 1989


Kjell Henriksen